

THE AFFIRMATIVE SOLUTION TO SALEM'S PROBLEM

SEMYON YAKUBOVICH

ABSTRACT. By using methods of classical analysis and special functions an old and attractive Salem's problem (posed in *Trans. Amer. Math. Soc.* **53** (3) (1943), p. 439) whether Fourier-Stieltjes coefficients of the Minkowski question mark function vanish at infinity is solved affirmatively. The coefficients d_n decay polynomially, namely, $d_n = O(n^{-\mu})$, $n \rightarrow \infty$ for some $\mu > 0$. Moreover, we generalize the Salem problem, proving that the Fourier-Stieltjes coefficients of any power $m \in \mathbb{N}$ of the Minkowski question mark function vanish at infinity as well.

Keywords: Minkowski question mark function, Salem's problem, Fourier-Stieltjes transform, Fourier-Stieltjes coefficients

Mathematics subject classification: 42A16, 42B10, 44A15

1. INTRODUCTION

Let $x \in \mathbb{R}$ and consider the following Fourier-Stieltjes transforms

$$f(x) = \int_0^1 e^{ixt} dq(t), \quad (1.1)$$

$$F(x) = \int_0^\infty e^{ixt} dq(t). \quad (1.2)$$

Here $q(x)$ is the famous Minkowski question mark function $?(x) \equiv q(x)$. This function is defined by [2] $q(x) : [0, 1] \mapsto [0, 1]$

$$q([0, a_1, a_2, a_3, \dots]) = 2 \sum_{i=1}^{\infty} (-1)^{i+1} 2^{-\sum_{j=1}^i a_j},$$

where $x = [0, a_1, a_2, a_3, \dots]$ stands for the representation of x by a regular continued fraction. It is well known that $q(x)$ is continuous, strictly increasing and supports a singular measure. It is uniquely determined by the following functional equations, which will be used in the sequel

$$q(x) = 1 - q(1 - x), \quad x \in [0, 1], \quad (1.3)$$

$$q(x) = 2q\left(\frac{x}{x+1}\right), \quad x \in [0, 1], \quad (1.4)$$

$$q(x) + q\left(\frac{1}{x}\right) = 2, \quad x > 0. \quad (1.5)$$

When $x \rightarrow 0$, it decreases exponentially $q(x) = O(2^{-1/x})$. Key values are $q(0) = 0$, $q(1) = 1$, $q(\infty) = 2$. For instance, from (1.3) and asymptotic behavior of the Minkowski function near

zero one can easily get the finiteness of the following integrals

$$\int_0^1 x^\lambda dq(x) < \infty, \lambda \in \mathbb{R}, \quad (1.6)$$

$$\int_0^1 (1-x)^\lambda dq(x) < \infty, \lambda \in \mathbb{R}. \quad (1.7)$$

Further, as was proved by Salem [9], the Minkowski question mark function satisfies the Hölder condition

$$|q(x) - q(y)| < C|x - y|^\alpha, \alpha < 1,$$

where

$$\alpha = \frac{\log 2}{2 \log \frac{\sqrt{5}+1}{2}} = 0,7202_+.$$

and $C > 0$ is an absolute constant. As we observe from the functional equation (1.3) the Fourier-Stieltjes transform (1.1) satisfies the functional relation

$$f(x) = e^{ix} f(-x), \quad (1.8)$$

and therefore $e^{-ix/2} f(x)$ is real-valued. So, taking its imaginary part, we obtain the equality

$$\cos\left(\frac{x}{2}\right) f_s(x) = \sin\left(\frac{x}{2}\right) f_c(x), \quad (1.9)$$

where f_s, f_c are the Fourier-Stieltjes sine and cosine transforms of the Minkowski question mark function, respectively,

$$f_s(x) = \int_0^1 \sin(xt) dq(t), \quad (1.10)$$

$$f_c(x) = \int_0^1 \cos(xt) dq(t). \quad (1.11)$$

Hence, letting, for instance, $x = 2\pi n$, $n \in \mathbb{N}_0$ it gives $f_s(2\pi n) = 0$ and $f_c(2\pi n) = d_n$. In 1943 Salem asked [9] whether $d_n \rightarrow 0$, as $n \rightarrow \infty$. This question is quite delicate, since it concerns singular functions (see [11], Ch. IV) and the classical Riemann-Lebesgue lemma for the class L_1 , in general, cannot be applied. A singular function is defined as a continuous, bounded monotone function with a null derivative almost everywhere. Hence it supports a positive bounded Borel measure, which is singular with respect to Lebesgue measure. For such singular measures there are various examples whose Fourier transforms do not tend to zero, although some do (see, for instance, in [9], [10], [5]). In [14] (see also [3]) it was proved that for every $\varepsilon > 0$ there exists a singular monotone function, which supports a measure whose Fourier-Stieltjes transform behaves as $O(t^{-\frac{1}{2}+\varepsilon})$, $|t| \rightarrow \infty$.

In fact, it is worth to mention that the Salem problem is an old and quite attractive problem in the number theory and Fourier analysis [13]. Several attempts were undertaken to solve Salem's problem (see, for instance, in [1], [16], [15]). Moreover, after appearance of the original version of this article [17] on Arxiv, it was noted in [7] that the solution to Salem's problem is a special case of a more general result of the paper [4].

In the sequel we will give the affirmative solution to Salem's problem, using the methods of classical analysis and special functions. To do this, we will employ all functional equations, which

uniquely describe the Minkowski question mark function and establish a new integro-differential equation for the Fourier-Stieltjes transform (1.1). It involves, in turn, the following functional equation, which is proved by the author in [16] and relates to transforms (1.1), (1.2)

$$f(x) = \left(1 - \frac{e^{ix}}{2}\right) F(x), \quad x \in \mathbb{R}. \quad (1.12)$$

Taking real and imaginary parts of both sides in (1.12), we derive interesting equalities (see details in [16]), which will be used below, namely

$$\int_1^\infty \cos xt \, dq(t) = \frac{1 - 8 \sin^2(x/2)}{1 + 8 \sin^2(x/2)} \int_0^1 \cos xt \, dq(t), \quad x \in \mathbb{R}, \quad (1.13)$$

$$\int_1^\infty \sin xt \, dq(t) = \frac{5 - 8 \sin^2(x/2)}{1 + 8 \sin^2(x/2)} \int_0^1 \sin xt \, dq(t), \quad x \in \mathbb{R}. \quad (1.14)$$

Making $x \rightarrow 0$ in (1.14), we find, in particular,

$$\int_1^\infty t \, dq(t) = 5 \int_0^1 t \, dq(t). \quad (1.15)$$

Moreover, using the functional equation (1.3), it can be proved the important equality for coefficients d_n

$$d_n = 2 \int_0^1 t \cos(2\pi nt) \, dq(t). \quad (1.16)$$

Indeed, we have

$$\begin{aligned} \int_0^1 t \cos(2\pi nt) \, dq(t) - \frac{d_n}{2} &= \left(\int_0^{1/2} + \int_{1/2}^1 \right) \left(t - \frac{1}{2} \right) \cos(2\pi nt) \, dq(t) \\ &= \int_0^{1/2} \left(t - \frac{1}{2} \right) \cos(2\pi nt) \, dq(t) - \int_{1/2}^1 \left(t - \frac{1}{2} \right) \cos(2\pi nt) \, dq(1-t) = 0. \end{aligned}$$

Further, we give values of the important relatively convergent integrals, which are calculated with the use of the Parseval equality for the Mellin transform [12] and verified recently with *Mathematica*. It will be employed in Section 3 to solve the Salem problem. Precisely, according to relations (2.5.22.2) and (2.5.22.6) in [8], Vol. 1 the following equalities hold for $a, b > 0$ and $0 < \mu < 2$

$$\begin{aligned} \int_0^\infty x^{\mu-1} \left\{ \frac{\sin(ax^2)}{\cos(ax^2)} \right\} \sin(bx) \, dx &= \frac{b}{2a^{(\mu+1)/2}} \Gamma\left(\frac{\mu+1}{2}\right) \left\{ \frac{\cos((1-\mu)\pi/4)}{\sin((1-\mu)\pi/4)} \right\} \\ \times {}_2F_3\left(\frac{\mu+3}{4}, \frac{\mu+1}{4}, \frac{1}{2}, \frac{3}{4}, \frac{5}{4}; -\left(\frac{b^2}{8a}\right)^2\right) &\mp \frac{b^3}{12a^{(\mu+3)/2}} \Gamma\left(\frac{\mu+3}{2}\right) \left\{ \frac{\cos((1+\mu)\pi/4)}{\sin((1+\mu)\pi/4)} \right\} \end{aligned}$$

$$\times {}_2F_3\left(\frac{\mu+5}{4}, \frac{\mu+3}{4}; \frac{3}{2}, \frac{5}{4}, \frac{7}{4}; -\left(\frac{b^2}{8a}\right)^2\right), \quad (1.17)$$

$$\begin{aligned} \int_0^\infty x^{\mu-1} \left\{ \frac{\sin(ax^2)}{\cos(ax^2)} \right\} \cos(bx) dx &= \frac{1}{2a^{\mu/2}} \Gamma\left(\frac{\mu}{2}\right) \left\{ \frac{\sin(\mu\pi/4)}{\cos(\mu\pi/4)} \right\} \\ &\times {}_2F_3\left(\frac{\mu+2}{4}, \frac{\mu}{4}; \frac{1}{2}, \frac{1}{4}, \frac{3}{4}; -\left(\frac{b^2}{8a}\right)^2\right) \mp \frac{b^2}{4a^{\mu/2+1}} \Gamma\left(\frac{\mu}{2}+1\right) \left\{ \frac{\cos(\mu\pi/4)}{\sin(\mu\pi/4)} \right\} \\ &\times {}_2F_3\left(\frac{\mu}{4}+1, \frac{\mu+2}{4}; \frac{3}{2}, \frac{3}{4}, \frac{5}{4}; -\left(\frac{b^2}{8a}\right)^2\right). \end{aligned} \quad (1.18)$$

Here $\Gamma(z)$ is Euler's gamma-function and ${}_2F_3(\alpha_1, \alpha_2; \beta_1, \beta_2, \beta_3; -x)$ is the generalized hypergeometric function, having the following asymptotic behavior at infinity, which is confirmed with *Mathematica* (see in [6], Section 16.11 (ii))

$$\begin{aligned} &{}_2F_3(\alpha_1, \alpha_2; \beta_1, \beta_2, \beta_3; -x) \\ &= \frac{\Gamma(\beta_1)\Gamma(\beta_2)\Gamma(\beta_3)}{\Gamma(\alpha_1)\Gamma(\alpha_2)} \left[\frac{x^\gamma}{\sqrt{\pi}} \cos(2\sqrt{x} + \pi\gamma) + O(x^{\gamma-1/2}) + O(x^{-\alpha_1}) + O(x^{-\alpha_2}) \right], \quad x \rightarrow +\infty, \end{aligned} \quad (1.19)$$

where

$$\gamma = \frac{1}{4} + \frac{1}{2} \left[\sum_{j=1}^2 \alpha_j - \sum_{j=1}^3 \beta_j \right]. \quad (1.20)$$

2. INTEGRO-DIFFERENTIAL EQUATION FOR THE FOURIER-STIELTJES TRANSFORM (1.1)

In order to make the paper self-contained we begin with the proof of the relation (1.12) (cf. [16]).

Lemma 1. *Let $x \in \mathbb{R}$ and $f(x)$, $F(x)$ be Fourier-Stieltjes transforms (1.1), (1.2), respectively. Then functional equation (1.12) holds.*

Proof. The proof is based on functional equations (1.4), (1.5) for the Minkowski question mark function and simple properties of the Stieltjes integral. In fact, we derive the chain of equalities

$$\begin{aligned}
\int_0^1 e^{ixt} dq(t) &= \int_0^\infty e^{ixt} dq(t) - \int_1^\infty e^{ixt} dq(t) \\
&= \int_0^\infty e^{ixt} dq(t) - e^{ix} \int_0^\infty e^{ixt} dq(t+1) \\
&= \int_0^\infty e^{ixt} dq(t) + e^{ix} \int_0^\infty e^{ixt} dq\left(\frac{1}{t+1}\right) \\
&= \int_0^\infty e^{ixt} dq(t) + e^{ix} \int_0^\infty e^{ixt} dq\left(\frac{1/t}{1+1/t}\right) \\
&= \int_0^\infty e^{ixt} dq(t) + \frac{e^{ix}}{2} \int_0^\infty e^{ixt} dq\left(\frac{1}{t}\right) \\
&= \left(1 - \frac{e^{ix}}{2}\right) \int_0^\infty e^{ixt} dq(t),
\end{aligned}$$

which yields (1.12). $\square$

Theorem 1. Let $x \in \mathbb{R}_+$. The Fourier-Stieltjes transform (1.1) satisfies the following integro-differential equation, involving the operator of the modified Hankel transform

$$\frac{e^{ix}}{2 - e^{ix}} \left[f'(x) + \frac{2if(x)}{2 - e^{ix}} \right] = - \int_0^\infty J_0(2\sqrt{xy}) e^{-iy} f(y) dy. \quad (2.1)$$

Proof. Indeed, differentiating (1.12) with respect to x and using it again, we find

$$f'(x) = -\frac{ie^{ix}f(x)}{2 - e^{ix}} + i\left(1 - \frac{e^{ix}}{2}\right) \int_0^\infty te^{ixt} dq(t), \quad (2.2)$$

where the differentiation under the integral sign in (1.2) is allowed via the simple estimate

$$\left| \int_0^\infty te^{ixt} dq(t) \right| \leq \int_0^\infty t dq(t) = 3,$$

where the latter equality is due to (1.15), (1.16). Hence,

$$\begin{aligned}
i \int_0^\infty t e^{ixt} dq(t) &= i \int_0^1 te^{ixt} dq(t) + i \int_1^\infty te^{ixt} dq(t) \\
&= f'(x) + i \int_0^1 \frac{e^{ix/t}}{t} dq(t).
\end{aligned} \quad (2.3)$$

Recalling the relatively convergent integral from [8], relation (2.12.9.3)

$$\frac{e^{ix/t}}{it} = \int_0^\infty J_0(2\sqrt{xy}) e^{-ity} dy, \quad x, t > 0, \quad (2.4)$$

where $J_0(z)$ is the Bessel function of the first kind [8], Vol. 2, we substitute it in (2.3). Hence after the change of the order of integration and the use of the symmetry property (1.8), we combine with (2.2) and come up with the integro-differential equation (2.1). Our goal now is to motivate the interchange of the order of integration in the iterated integral, proving the formula

$$\int_0^1 \left(\int_0^\infty J_0(2\sqrt{xy}) e^{-ity} dy \right) dq(t) = \int_0^\infty J_0(2\sqrt{xy}) \left(\int_0^1 e^{-ity} dq(t) \right) dy, \quad x > 0. \quad (2.5)$$

To do this, it is sufficient to justify the limit equality

$$\lim_{Y \rightarrow \infty} \int_0^1 \left(\int_Y^\infty J_0(2\sqrt{xy}) e^{-ity} dy \right) dq(t) = 0 \quad (2.6)$$

for each fixed positive x . Naturally, we will appeal to the known asymptotic behavior of the Bessel function at infinity [6], Section 10.17 (i)

$$J_\nu(y) = \sqrt{\frac{2}{\pi y}} \left[\cos\left(y - \frac{\pi\nu}{2} - \frac{\pi}{4}\right) - \frac{a(\nu)}{y} \sin\left(y - \frac{\pi\nu}{2} - \frac{\pi}{4}\right) + O\left(\frac{1}{y^2}\right) \right], \quad y \rightarrow +\infty, \quad (2.7)$$

where

$$a(\nu) = \frac{\nu^2}{2} - \frac{1}{8}, \quad \nu \in \mathbb{R}.$$

Hence, for sufficiently large $Y > 0$ and $x > 0$, $t \in (0, 1)$, we have

$$\begin{aligned} \int_Y^\infty J_0(2\sqrt{xy}) e^{-ity} dy &= \frac{1}{\sqrt{\pi x^{1/4}}} \int_Y^\infty \cos\left(2\sqrt{xy} - \frac{\pi}{4}\right) e^{-ity} \frac{dy}{y^{1/4}} \\ &+ \frac{1}{16\sqrt{\pi x^{3/4}}} \int_Y^\infty \sin\left(2\sqrt{xy} - \frac{\pi}{4}\right) e^{-ity} \frac{dy}{y^{3/4}} + O\left(Y^{-1/4}\right). \end{aligned} \quad (2.8)$$

As we will see from the estimates below and the finiteness of integrals (1.6) for various real λ , in order to establish the limit (2.6), it is sufficient to estimate, for instance, the integral

$$\int_Y^\infty \cos(2\sqrt{xy}) \cos(ty) \frac{dy}{y^{1/4}},$$

because other integrals in (2.8) can be estimated in the same manner. With the simple substitution and integration by parts we have

$$\begin{aligned} \int_Y^\infty \cos(2\sqrt{xy}) \cos(ty) \frac{dy}{y^{1/4}} &= 2 \int_{\sqrt{Y}}^\infty \cos(2y\sqrt{x}) \cos(ty^2) \sqrt{y} dy \\ &= -\frac{\cos(2\sqrt{xY}) \sin(tY)}{tY^{1/4}} + \frac{1}{2t} \int_{\sqrt{Y}}^\infty \sin(ty^2) \left[\frac{\cos(2y\sqrt{x})}{y} + 4\sqrt{x} \sin(2y\sqrt{x}) \right] \frac{dy}{\sqrt{y}} \\ &= \frac{2\sqrt{x}}{t} \int_{\sqrt{Y}}^\infty \sin(ty^2) \sin(2y\sqrt{x}) \frac{dy}{\sqrt{y}} + O\left(t^{-1}Y^{-1/4}\right). \end{aligned}$$

Similarly,

$$\begin{aligned} \frac{2\sqrt{x}}{t} \int_{\sqrt{Y}}^\infty \sin(ty^2) \sin(2y\sqrt{x}) \frac{dy}{\sqrt{y}} &= O\left(t^{-2}Y^{-3/4}\right) \\ + \frac{\sqrt{x}}{2t^2} \int_{\sqrt{Y}}^\infty \cos(ty^2) \left[-\frac{3\sin(2y\sqrt{x})}{y} + 4\sqrt{x} \cos(2y\sqrt{x}) \right] \frac{dy}{y^{3/2}} \end{aligned}$$

$$= O\left(t^{-2}Y^{-1/4}\right).$$

Consequently,

$$\begin{aligned} & \int_0^1 \left(\int_Y^\infty \cos(2\sqrt{xy}) \cos(ty) \frac{dy}{y^{1/4}} \right) dq(t) \\ &= O\left(Y^{-1/4} \left[\int_0^1 t^{-1} dq(t) + \int_0^1 t^{-2} dq(t) \right] \right) = O\left(Y^{-1/4}\right), Y \rightarrow \infty. \end{aligned}$$

Therefore, treating in the same manner other integrals from (2.8), we get equality (2.6), completing the proof of Theorem 1. $\square$

Remark 1. A similar to (2.1) integro-differential equation for the Fourier-Stieltjes transform (1.1) with the derivative $f'(x)$ inside the modified Hankel transform [12] was exhibited in [1]. However it did not lead to the solution of the Salem problem.

Corollary 1. Let $n \in \mathbb{N}$. The values $d_n = f(2\pi n)$ and

$$c_n = \int_0^1 t \sin(2\pi nt) dq(t) \quad (2.9)$$

have the following integral representations in terms of the modified Hankel transform

$$d_n = \frac{2}{5} \int_0^\infty J_0\left(2\sqrt{2\pi ny}\right) f_s(y) dy, \quad (2.10)$$

$$c_n = \int_0^\infty J_0\left(2\sqrt{2\pi ny}\right) f_c(y) dy, \quad (2.11)$$

where $f_s(x), f_c(x)$ are the Fourier-Stieltjes sine and cosine transforms of the Minkowski question mark function (1.10), (1.11), respectively.

Proof. Indeed, substituting in (2.1) $x = 2\pi n$, we have

$$f'(2\pi n) + 2id_n = - \int_0^\infty J_0\left(2\sqrt{2\pi ny}\right) e^{-iy} f(y) dy.$$

In the meantime, it is not difficult to show, recalling (1.16), that

$$\begin{aligned} f'(2\pi n) &= i \int_0^1 t e^{2\pi i nt} dq(t) = i \int_0^1 t \cos(2\pi nt) dq(t) \\ &\quad - \int_0^1 t \sin(2\pi nt) dq(t) = \frac{i}{2} d_n - c_n. \end{aligned} \quad (2.12)$$

Hence,

$$\frac{5}{2} id_n - c_n = - \int_0^\infty J_0\left(2\sqrt{2\pi ny}\right) e^{-iy} f(y) dy. \quad (2.13)$$

Now taking the imaginary and real parts of both sides of the latter equality in (2.13) with the use of (1.3), we end up with (2.10), (2.11). $\square$

3. SOLUTION TO SALEM'S PROBLEM

The main result is the following

Theorem 2. *The answer on Salem's question is affirmative. Moreover, there exists some $\mu > 0$ such that*

$$d_n = O(n^{-\mu}), \quad n \rightarrow \infty.$$

Proof. Indeed, taking (2.10), we write

$$\frac{5}{2}d_n = \int_0^1 J_0(2\sqrt{2\pi ny}) f_s(y) dy + \int_1^\infty J_0(2\sqrt{2\pi ny}) f_s(y) dy = I_1(n) + I_2(n). \quad (3.1)$$

From the inequality $\sqrt{x}|J_\nu(x)| \leq C$, $x > 0$, where $C > 0$ is an absolute constant, it follows that $I_1(n)$ converges absolutely and uniformly with respect to n . Therefore $I_1(n) \rightarrow 0$, $n \rightarrow \infty$. Concerning the integral $I_2(n)$, we appeal to the asymptotic formula (2.7) for the Bessel function and recall (1.10) to get the equalities

$$\begin{aligned} I_2(n) &= \int_1^\infty J_0(2\sqrt{2\pi ny}) f_s(y) dy \\ &= \frac{1}{\sqrt{\pi}(2\pi n)^{1/4}} \int_1^\infty \cos\left(2\sqrt{2\pi ny} - \frac{\pi}{4}\right) \left(\int_0^1 \sin(yt) dq(t)\right) \frac{dy}{y^{1/4}} \\ &\quad + \frac{1}{16\sqrt{\pi}(2\pi n)^{3/4}} \int_1^\infty \sin\left(2\sqrt{2\pi ny} - \frac{\pi}{4}\right) \left(\int_0^1 \sin(yt) dq(t)\right) \frac{dy}{y^{3/4}} \\ &\quad + O\left(\frac{1}{n^{5/4}} \int_1^\infty \left(\int_0^1 \sin(yt) dq(t)\right) \frac{dy}{y^{5/4}}\right) = I_{21}(n) + I_{22}(n) + I_{23}(n). \end{aligned} \quad (3.2)$$

Hence

$$|I_{23}(n)| \leq \frac{C}{n^{5/4}} \int_0^1 dq(t) = O(n^{-5/4}), \quad n \rightarrow \infty,$$

where $C > 0$ is an absolute constant. Further, going backwards by the interchange of the order of integration in the integral I_{22} for each n as in the proof of Theorem 1, we write it in the form

$$\begin{aligned} I_{22}(n) &= \frac{1}{16\sqrt{\pi}(2\pi n)^{3/4}} \int_1^\infty \sin\left(2\sqrt{2\pi ny} - \frac{\pi}{4}\right) \left(\int_0^1 \sin(yt) dq(t)\right) \frac{dy}{y^{3/4}} \\ &= \frac{1}{16(2\pi)^{5/4}n^{3/4}} \int_0^1 dq(t) \left(\int_0^\infty - \int_0^1\right) \left[\sin\left(2\sqrt{2\pi ny}\right) - \cos\left(2\sqrt{2\pi ny}\right)\right] \sin(yt) \frac{dy}{y^{3/4}}, \end{aligned} \quad (3.3)$$

where by virtue of the absolute and uniform convergence with respect to $n \in \mathbb{N}$ and $t \in [0, 1]$

$$\begin{aligned} &\frac{1}{16(2\pi)^{5/4}n^{3/4}} \int_0^1 \left[\sin\left(2\sqrt{2\pi ny}\right) - \cos\left(2\sqrt{2\pi ny}\right)\right] \sin(yt) \frac{dy}{y^{3/4}} \\ &= O(n^{-3/4}), \quad n \rightarrow \infty. \end{aligned} \quad (3.4)$$

In the mean time, the corresponding relatively convergent integral (3.4) over $(0, \infty)$ is calculated via equalities (1.17), (1.18), letting there $a = t/(2\pi n)$, $b = 2$, $\mu = 1/2$. In fact, making an elementary substitution, we find

$$\frac{1}{16(2\pi)^{5/4}n^{3/4}} \int_0^\infty \left[\sin\left(2\sqrt{2\pi ny}\right) - \cos\left(2\sqrt{2\pi ny}\right)\right] \sin(yt) \frac{dy}{y^{3/4}}$$

$$\begin{aligned}
&= \frac{1}{8(2\pi)^{3/2}n} \int_0^\infty [\sin(2y) - \cos(2y)] \sin\left(y^2 \frac{t}{2\pi n}\right) \frac{dy}{y^{1/2}} \\
&= \frac{1}{8(2\pi)^{3/4}n^{1/4}t^{3/4}} \Gamma\left(\frac{3}{4}\right) \cos\left(\frac{\pi}{8}\right) {}_2F_3\left(\frac{7}{8}, \frac{3}{8}; \frac{1}{2}, \frac{3}{4}, \frac{5}{4}; -\left(\frac{\pi n}{t}\right)^2\right) \\
&\quad - \frac{(2\pi)^{1/4}n^{3/4}}{12t^{7/4}} \Gamma\left(\frac{7}{4}\right) \cos\left(\frac{3\pi}{8}\right) {}_2F_3\left(\frac{11}{8}, \frac{7}{8}; \frac{3}{2}, \frac{5}{4}, \frac{7}{4}; -\left(\frac{\pi n}{t}\right)^2\right) \\
&\quad - \frac{1}{16(2\pi)^{5/4}n^{3/4}t^{1/4}} \Gamma\left(\frac{1}{4}\right) \sin\left(\frac{\pi}{8}\right) {}_2F_3\left(\frac{5}{8}, \frac{1}{8}; \frac{1}{2}, \frac{1}{4}, \frac{3}{4}; -\left(\frac{\pi n}{t}\right)^2\right) \\
&\quad + \frac{n^{1/4}}{8(2\pi)^{1/4}t^{5/4}} \Gamma\left(\frac{5}{4}\right) \cos\left(\frac{\pi}{8}\right) {}_2F_3\left(\frac{9}{8}, \frac{5}{8}; \frac{3}{2}, \frac{3}{4}, \frac{5}{4}; -\left(\frac{\pi n}{t}\right)^2\right). \tag{3.5}
\end{aligned}$$

Hence, appealing to the asymptotic formula (1.19) and calculating the corresponding parameter γ by formula (1.20), we establish the asymptotic behavior of the right-hand side of the latter equality in (3.5) when $n \rightarrow \infty$ and $t \in (0, 1)$. In fact, taking the contribution of each hypergeometric function, we obtain

$$\begin{aligned}
{}_2F_3\left(\frac{7}{8}, \frac{3}{8}; \frac{1}{2}, \frac{3}{4}, \frac{5}{4}; -\left(\frac{\pi n}{t}\right)^2\right) &= \frac{\Gamma(3/4)\Gamma(5/4)}{\Gamma(7/8)\Gamma(3/8)} \left(\frac{\pi n}{t}\right)^{-3/4} \cos\left(\frac{2\pi n}{t} - \frac{3\pi}{8}\right) \\
&\quad + O\left(\left(\frac{n}{t}\right)^{-3/4}\right), \tag{3.6}
\end{aligned}$$

$$\begin{aligned}
{}_2F_3\left(\frac{11}{8}, \frac{7}{8}; \frac{3}{2}, \frac{5}{4}, \frac{7}{4}; -\left(\frac{\pi n}{t}\right)^2\right) &= \frac{\Gamma(5/4)\Gamma(7/4)}{2\Gamma(11/8)\Gamma(7/8)} \left(\frac{\pi n}{t}\right)^{-7/4} \cos\left(\frac{2\pi n}{t} - \frac{7\pi}{8}\right) \\
&\quad + O\left(\left(\frac{n}{t}\right)^{-7/4}\right), \tag{3.7}
\end{aligned}$$

$$\begin{aligned}
{}_2F_3\left(\frac{5}{8}, \frac{1}{8}; \frac{1}{2}, \frac{1}{4}, \frac{3}{4}; -\left(\frac{\pi n}{t}\right)^2\right) &= \frac{\Gamma(1/4)\Gamma(3/4)}{\Gamma(5/8)\Gamma(1/8)} \left(\frac{\pi n}{t}\right)^{-1/4} \cos\left(\frac{2\pi n}{t} - \frac{\pi}{8}\right) \\
&\quad + O\left(\left(\frac{n}{t}\right)^{-1/4}\right), \tag{3.8}
\end{aligned}$$

$$\begin{aligned}
{}_2F_3\left(\frac{9}{8}, \frac{5}{8}; \frac{3}{2}, \frac{3}{4}, \frac{5}{4}; -\left(\frac{\pi n}{t}\right)^2\right) &= \frac{\Gamma(3/4)\Gamma(5/4)}{2\Gamma(9/8)\Gamma(5/8)} \left(\frac{\pi n}{t}\right)^{-5/4} \cos\left(\frac{2\pi n}{t} - \frac{5\pi}{8}\right) \\
&\quad + O\left(\left(\frac{n}{t}\right)^{-5/4}\right). \tag{3.9}
\end{aligned}$$

Hence, combining with (3.5) it is not difficult to see that its left-hand side is $O(n^{-1})$, $n \rightarrow \infty$ uniformly by $t \in (0, 1)$. Therefore, recalling (3.3), (3.4), we find the estimate

$$I_{22}(n) = O\left(n^{-3/4}\right), \quad n \rightarrow \infty. \tag{3.10}$$

The main obstacle is to estimate the integral $I_{21}(n)$ in (3.2). To do this we represent it in the form

$$I_{21}(n) = \frac{1}{\sqrt{\pi}(2\pi n)^{1/4}} \int_1^\infty \cos\left(2\sqrt{2\pi n y} - \frac{\pi}{4}\right) \left(\int_0^1 \sin(yt) dq(t)\right) \frac{dy}{y^{1/4}}$$

$$\begin{aligned}
&= \frac{1}{2\sqrt{\pi}(2\pi n)^{1/4}} \int_0^1 dq(t) \int_1^\infty \sin\left(2\sqrt{2\pi n y} + yt - \frac{\pi}{4}\right) \frac{dy}{y^{1/4}} \\
&\quad - \frac{1}{2\sqrt{\pi}(2\pi n)^{1/4}} \int_0^1 dq(t) \int_1^\infty \sin\left(2\sqrt{2\pi n y} - yt - \frac{\pi}{4}\right) \frac{dy}{y^{1/4}}.
\end{aligned} \tag{3.11}$$

But the derivative of the function $h(y) = 2\sqrt{2\pi n y} + yt - \frac{\pi}{4}$

$$h'(y) = \sqrt{\frac{2\pi n}{y}} + t, \quad y \geq 1, \quad t \in (0, 1)$$

is monotone decreasing. Therefore via lemma of Section 1.12 in [12]

$$\int_1^\infty \sin\left(2\sqrt{2\pi n y} + yt - \frac{\pi}{4}\right) \frac{dy}{y^{1/4}} = O\left(\max_{y \geq 1} \left[\sqrt{\frac{2\pi n}{y}} + t\right]^{-1}\right) = O\left(\frac{1}{t}\right).$$

Consequently,

$$\frac{1}{2\sqrt{\pi}(2\pi n)^{1/4}} \int_0^1 dq(t) \int_1^\infty \sin\left(2\sqrt{2\pi n y} + yt - \frac{\pi}{4}\right) \frac{dy}{y^{1/4}} = O(n^{-1/4}), \quad n \rightarrow \infty.$$

Returning to (3.11) and employing (1.8), we estimate the integral

$$\begin{aligned}
&-\frac{1}{2\sqrt{\pi}(2\pi n)^{1/4}} \int_0^1 dq(t) \int_1^\infty \sin\left(2\sqrt{2\pi n y} - yt - \frac{\pi}{4}\right) \frac{dy}{y^{1/4}} \\
&= -\frac{1}{4i\sqrt{\pi}(2\pi n)^{1/4}} \int_0^1 dq(t) \int_1^\infty \exp\left(i\left(2\sqrt{2\pi n y} - y(1-t) - \frac{\pi}{4}\right)\right) \frac{dy}{y^{1/4}} \\
&\quad + \frac{1}{4i\sqrt{\pi}(2\pi n)^{1/4}} \int_0^1 dq(t) \int_1^\infty \exp\left(i\left(\frac{\pi}{4} + y(1-t) - 2\sqrt{2\pi n y}\right)\right) \frac{dy}{y^{1/4}}.
\end{aligned} \tag{3.12}$$

Let us examine the first inner integral with respect to y in the right-hand side of the equality (3.12). Indeed, the derivative by y of the function

$$\varphi_n(y, t) = 2\sqrt{2\pi n y} - y(1-t) - \frac{\pi}{4}$$

vanishes at the point

$$y_n(t) = \frac{2\pi n}{(1-t)^2}, \quad t \in (0, 1), \quad n \in \mathbb{N}$$

and the second derivative

$$\varphi_n''(y_n) = -\frac{(1-t)^3}{4\pi n}.$$

Let

$$\rho(n) = n^\xi, \quad \frac{1}{2} < \xi < \frac{5}{8}. \tag{3.13}$$

Then for any $u \in \mathbb{R}$ such that $|u| \leq \rho(n)$ and n is big enough

$$|\varphi_n''(y_n + u, t) - \varphi_n''(y_n, t)| = \frac{(1-t)^3}{4\pi n} \left[1 - \left(1 + \frac{u(1-t)^2}{2\pi n}\right)^{-3/2}\right] < A (1-t)^5 n^{\xi-2}, \tag{3.14}$$

where $0 \leq u \leq \rho(n)$ and $A > 0$ is an absolute constant,

$$|\varphi_n''(y_n + u, t) - \varphi_n''(y_n, t)| \leq \frac{(1-t)^3}{4\pi n} \left[\left(1 - \frac{|u|(1-t)^2}{2\pi n} \right)^{-3/2} - 1 \right] < B (1-t)^5 n^{\xi-2}, \quad (3.15)$$

where $-\rho(n) \leq u \leq 0$ and $B > 0$ is an absolute constant. Now splitting the first inner integral with respect to y in the right-hand side of (3.12) (say, $G_n^+(t)$) on three integrals, namely,

$$\begin{aligned} G_n^+(t) &= -\frac{1}{4i\sqrt{\pi}(2\pi n)^{1/4}} \int_1^\infty \exp(i\varphi_n(y, t)) \frac{dy}{y^{1/4}} = -\frac{1}{4i\sqrt{\pi}(2\pi n)^{1/4}} \left[\int_1^{-\rho(n)+y_n} \exp(i\varphi_n(y, t)) \frac{dy}{y^{1/4}} \right. \\ &\quad \left. + \int_{-\rho(n)+y_n}^{\rho(n)+y_n} \exp(i\varphi_n(y, t)) \frac{dy}{y^{1/4}} + \int_{\rho(n)+y_n}^\infty \exp(i\varphi_n(y, t)) \frac{dy}{y^{1/4}} \right] = G_{1n}^+(t) + G_{2n}^+(t) + G_{3n}^+(t), \quad (3.16) \end{aligned}$$

we estimate each of them separately. In fact, integration by parts gives

$$\begin{aligned} -4i\sqrt{\pi}(2\pi)^{1/4} G_{1n}^+(t) &= -i (ny)^{-1/4} \frac{\exp(i\varphi_n(y, t))}{\varphi_n'(y, t)} \Big|_1^{-\rho(n)+y_n} - \frac{i}{4n^{1/4}} \int_1^{-\rho(n)+y_n} \frac{\exp(i\varphi_n(y, t))}{\varphi_n'(y, t) y^{5/4}} dy \\ &\quad - \frac{i}{n^{1/4}} \int_1^{-\rho(n)+y_n} \frac{\exp(i\varphi_n(y, t)) \varphi_n''(y, t)}{(\varphi_n'(y, t))^2 y^{1/4}} dy. \end{aligned}$$

Hence, taking into account the integrated terms and the value of the second derivative

$$\varphi_n''(y, t) = -\sqrt{\frac{\pi n}{2y^3}},$$

we end up with the uniform estimates

$$\begin{aligned} \frac{\exp(i\varphi_n(y, t))}{y^{1/4} \varphi_n'(y, t)} \Big|_1^{-\rho(n)+y_n} &= O(n^{-1/2}) + O((1-t)^{-5/2} n^{3/4-\xi}), \\ \left| \int_1^{-\rho(n)+y_n} \frac{\exp(i\varphi_n(y, t))}{\varphi_n'(y, t) y^{5/4}} dy \right| &\leq (2\pi n)^{-1/2} \int_1^{-\rho(n)+y_n} \left[1 - (1-t) \sqrt{\frac{y}{2\pi n}} \right]^{-1} \frac{dy}{y^{3/4}} \\ &= O((1-t)^{-2} n^{1/2-\xi}) + O((1-t)^{-5/2} n^{3/4-\xi}). \\ \left| \int_1^{-\rho(n)+y_n} \frac{\exp(i\varphi_n(y, t)) \varphi_n''(y, t)}{(\varphi_n'(y, t))^2 y^{1/4}} dy \right| &\leq - \int_1^{-\rho(n)+y_n} \frac{\varphi_n''(y, t)}{(\varphi_n'(y, t))^2} dy \\ &= O(n^{-1/2}) + O((1-t)^{-2} n^{1/2-\xi}). \end{aligned}$$

Returning to (3.16), we find, correspondingly, the estimate of $G_{1n}^+(t)$. Hence, calling (1.7) and combining with (3.12), we have

$$\int_0^1 G_{1n}^+(t) dq(t) = O(n^{-3/4}) + O(n^{1/2-\xi} \int_0^1 (1-t)^{-5/2} dq(t))$$

$$+O\left(n^{1/4-\xi}\int_0^1(1-t)^{-2}dq(t)\right)=O(n^{-\mu}),\ \mu>0,\ n\rightarrow\infty \quad (3.17)$$

because $\xi \in (1/2, 5/8)$ (see (3.13)).

In the meantime, for sufficiently big n and $y \in [y_n + \rho(n), \infty)$

$$\begin{aligned} |\varphi'_n(y, t)| &= 1 - t - \sqrt{\frac{2\pi n}{y}} \geq 1 - t - \sqrt{\frac{2\pi n}{y_n + \rho(n)}} \\ &= \frac{(1-t)^3 n^\xi}{2\pi n + (1-t)^2 n^\xi} \left(\sqrt{\frac{2\pi n}{2\pi n + (1-t)^2 n^\xi}} + 1 \right)^{-1} \geq C(1-t)^3 n^{\xi-1}, \end{aligned}$$

where $C > 0$ is an absolute constant. Then we do a similar analysis with the third integral

$$\begin{aligned} -4i\sqrt{\pi}(2\pi)^{1/4}G_{3n}^+(t) &= \int_{\rho(n)+y_n}^{\infty} \exp(i\varphi_n(y, t)) \frac{dy}{(ny)^{1/4}} \\ &= i \frac{\exp(i\varphi_n(y_n + \rho(n), t))}{(n(\rho(n) + y_n))^{1/4} \varphi'_n(y_n + \rho(n), t)} - \frac{i}{4n^{1/4}} \int_{\rho(n)+y_n}^{\infty} \frac{\exp(i\varphi_n(y, t))}{\varphi'_n(y, t) y^{5/4}} dy \\ &\quad - \frac{i}{n^{1/4}} \int_{\rho(n)+y_n}^{\infty} \frac{\exp(i\varphi_n(y, t)) \varphi''_n(y, t)}{(\varphi'_n(y, t))^2 y^{1/4}} dy = O\left((1-t)^{-5/2} n^{1/2-\xi}\right) \end{aligned}$$

and, correspondingly,

$$\int_0^1 G_{3n}^+(t) dq(t) = O(n^{-\mu}),\ \mu > 0,\ n \rightarrow \infty. \quad (3.18)$$

Concerning the integral $G_{2n}^+(t)$, we have by virtue of the Taylor formula when $y \in [y_n - \rho(n), y_n + \rho(n)]$

$$\varphi_n(y, t) = \varphi_n(y_n, t) + \frac{1}{2} \varphi''_n(h(y, n, t)) (y - y_n)^2,$$

where $|y_n(t) - h(y, n, t)| \leq \rho(n)$ for all $t \in (0, 1)$. Hence using the elementary inequality $|1 - e^{i\theta}| \leq |\theta|$, $\theta \in \mathbb{R}$, estimates (3.14), (3.15) and elementary substitutions, we write

$$\begin{aligned} -4i\sqrt{\pi}(2\pi)^{1/4}G_{2n}^+(t) &= \int_{-\rho(n)+y_n}^{\rho(n)+y_n} \exp(i\varphi_n(y, t)) \frac{dy}{(ny)^{1/4}} \\ &= \frac{\exp(i\varphi_n(y_n, t))}{n^{1/4}} \int_{-\rho(n)}^{\rho(n)} \left[\exp\left(\frac{i}{2} \varphi''_n(h(y, n, t)) y^2\right) - \exp\left(\frac{i}{2} \varphi''_n(y_n, t) y^2\right) \right] \frac{dy}{(y + y_n)^{1/4}} \\ &\quad + \frac{\exp(i\varphi_n(y_n, t))}{n^{1/4}} \int_{-\rho(n)}^{\rho(n)} \exp\left(\frac{i}{2} \varphi''_n(y_n, t) y^2\right) \frac{dy}{(y + y_n)^{1/4}} \end{aligned}$$

and since

$$\begin{aligned}
& \left| \int_{-\rho(n)}^{\rho(n)} \left[\exp \left(\frac{i}{2} \varphi_n''(h(y, n, t)) y^2 \right) - \exp \left(\frac{i}{2} \varphi_n''(y_n, t) y^2 \right) \right] \frac{dy}{(y + y_n)^{1/4}} \right| \\
& \leq \int_{-\rho(n)}^{\rho(n)} \left| \exp \left(\frac{i}{2} [\varphi_n''(h(y, n, t)) - \varphi_n''(y_n, t)] y^2 \right) - 1 \right| \frac{dy}{(y + y_n)^{1/4}} \leq A(1-t)^{5,5} n^{4\xi-9/4},
\end{aligned}$$

where $A > 0$ is an absolute constant, it gives via (3.13), (1.3)

$$\begin{aligned}
& \int_0^1 G_{2n}^+(t) dq(t) = -\frac{1}{4i\sqrt{\pi}(2\pi n)^{1/4}} \int_0^1 \int_{-n^\xi}^{n^\xi} \frac{\sqrt{t}}{(t^2 y + 2\pi n)^{1/4}} \\
& \times \exp \left(-i \left[\frac{t^3 y^2}{8\pi n} - \frac{2\pi n}{t} + \frac{\pi}{4} \right] \right) dy dq(t) + O(n^{-\mu}), \mu > 0, n \rightarrow \infty.
\end{aligned} \tag{3.19}$$

Meanwhile, with the use of the second mean value theorem there exists $\eta \in [-n^\xi, n^\xi]$ such that

$$\begin{aligned}
& \int_{-n^\xi}^{n^\xi} \left[\frac{1}{(t^2 y + 2\pi n)^{1/4}} - \frac{1}{(2\pi n)^{1/4}} \right] \exp \left(-\frac{t^3 y^2 i}{8\pi n} \right) dy = - \left[(2\pi n)^{1/4} (2\pi n - t^2 n^\xi)^{1/4} \left((2\pi n - t^2 n^\xi)^{1/4} \right. \right. \\
& \left. \left. + (2\pi n)^{1/4} \right) \left((2\pi n - t^2 n^\xi)^{1/2} + (2\pi n)^{1/2} \right) \right]^{-1} \int_{-n^\xi}^{\eta} t^2 y \exp \left(-\frac{t^3 y^2 i}{8\pi n} \right) dy = O \left(\frac{1}{t n^{1/4}} \right)
\end{aligned}$$

uniformly by $t \in (0, 1)$. Hence, minding (3.17), (3.18), we write the first iterated integral in the right-hand side of (3.12) as

$$\begin{aligned}
& \int_0^1 G_n^+(t) dq(t) = -\frac{n^{\xi-1/2}}{2i\sqrt{\pi}(2\pi)^{1/2}} \int_0^1 \int_0^1 \sqrt{t} \exp \left(-i \left[\frac{t^3 n^{2\xi-1} y^2}{8\pi} - \frac{2\pi n}{t} + \frac{\pi}{4} \right] \right) dy dq(t) \\
& + O(n^{-\mu}), n \rightarrow \infty.
\end{aligned} \tag{3.20}$$

Now, estimating analogously the integral

$$G_n^-(t) = \frac{1}{4i\sqrt{\pi}(2\pi n)^{1/4}} \int_1^\infty \exp(-i\varphi_n(y, t)) \frac{dy}{y^{1/4}},$$

the second iterated integral in the right-hand side of (3.12) becomes

$$\begin{aligned}
& \int_0^1 G_n^-(t) dq(t) = \frac{n^{\xi-1/2}}{2i\sqrt{\pi}(2\pi)^{1/2}} \int_0^1 \int_0^1 \sqrt{t} \exp \left(i \left[\frac{t^3 n^{2\xi-1} y^2}{8\pi} - \frac{2\pi n}{t} + \frac{\pi}{4} \right] \right) dy dq(t) \\
& + O(n^{-\mu}), n \rightarrow \infty.
\end{aligned} \tag{3.21}$$

Thus taking into account (3.1), (3.2), (3.4), (3.10), the latter equalities (3.20), (3.21) drive us at the following asymptotic equality

$$\frac{5}{2}d_n = \frac{n^{\xi-1/2}}{\pi\sqrt{2}} \int_0^1 \int_0^1 \sqrt{t} \sin \left(\frac{t^3 n^{2\xi-1} y^2}{8\pi} - \frac{2\pi n}{t} + \frac{\pi}{4} \right) dy dq(t) + O(n^{-\mu}), \quad n \rightarrow \infty. \quad (3.22)$$

Exactly the same analysis can be done, concerning equation (2.11), involving Fourier coefficients (2.9). We omit the details, writing the final asymptotic equality

$$c_n = \frac{n^{\xi-1/2}}{\pi\sqrt{2}} \int_0^1 \int_0^1 \sqrt{t} \cos \left(\frac{t^3 n^{2\xi-1} y^2}{8\pi} - \frac{2\pi n}{t} + \frac{\pi}{4} \right) dy dq(t) + O(n^{-\mu}), \quad n \rightarrow \infty. \quad (3.23)$$

Hence,

$$\frac{5}{2}d_n + c_n = \frac{n^{\xi-1/2}}{\pi} \int_0^1 \int_0^1 \sqrt{t} \cos \left(\frac{t^3 n^{2\xi-1} y^2}{8\pi} - \frac{2\pi n}{t} \right) dy dq(t) + O(n^{-\mu}), \quad n \rightarrow \infty, \quad (3.24)$$

$$\frac{5}{2}d_n - c_n = \frac{n^{\xi-1/2}}{\pi} \int_0^1 \int_0^1 \sqrt{t} \sin \left(\frac{t^3 n^{2\xi-1} y^2}{8\pi} - \frac{2\pi n}{t} \right) dy dq(t) + O(n^{-\mu}), \quad n \rightarrow \infty. \quad (3.25)$$

Writing (3.24) in the form

$$\begin{aligned} \frac{5}{2}d_n + c_n &= \frac{2\sqrt{2}}{\sqrt{\pi}} \int_0^1 \cos \left(\frac{2\pi n}{t} \right) \frac{1}{t} \int_0^{t^{3/2} n^{\xi-1/2}/\sqrt{8\pi}} \cos y^2 dy dq(t) \\ &+ \frac{2\sqrt{2}}{\sqrt{\pi}} \int_0^1 \sin \left(\frac{2\pi n}{t} \right) \frac{1}{t} \int_0^{t^{3/2} n^{\xi-1/2}/\sqrt{8\pi}} \sin y^2 dy dq(t) + O(n^{-\mu}), \end{aligned}$$

using the fundamental trigonometric identity and observing that

$$\int_0^\infty \cos y^2 dy = \int_0^\infty \sin y^2 dy = \frac{1}{2} \sqrt{\frac{\pi}{2}}, \quad (3.26)$$

$$\begin{aligned} \left| \int_{t^{3/2} n^{\xi-1/2}/\sqrt{8\pi}}^\infty \begin{Bmatrix} \cos y^2 \\ \sin y^2 \end{Bmatrix} dy \right| &= \frac{1}{2} \left| \int_{t^{3/2} n^{\xi-1/2}/\sqrt{8\pi}}^\infty \begin{Bmatrix} \cos y \\ \sin y \end{Bmatrix} \frac{dy}{\sqrt{y}} \right| \\ &= \frac{n^{1/2-\xi} \sqrt{8\pi}}{2 t^{3/2}} \left| \int_{t^{3/2} n^{\xi-1/2}/\sqrt{8\pi}}^T \begin{Bmatrix} \cos y \\ \sin y \end{Bmatrix} dy \right| \leq \sqrt{\frac{8\pi}{t^3}} n^{1/2-\xi}, \end{aligned}$$

we have

$$\begin{aligned} 5d_n + 2c_n &= \frac{n^{\xi-1/2}}{\pi} \int_0^1 \cos \left(\frac{2\pi n}{t} \right) \sqrt{t} dq(t) + \frac{n^{\xi-1/2}}{\pi} \int_0^1 \sin \left(\frac{2\pi n}{t} \right) \sqrt{t} dq(t) \\ &+ O(n^{-\mu}), \quad n \rightarrow \infty. \end{aligned}$$

Analogously, from (3.25) we find

$$5d_n - 2c_n = \frac{n^{\xi-1/2}}{\pi} \int_0^1 \cos\left(\frac{2\pi n}{t}\right) \sqrt{t} dq(t) - \frac{n^{\xi-1/2}}{\pi} \int_0^1 \sin\left(\frac{2\pi n}{t}\right) \sqrt{t} dq(t) \\ + O(n^{-\mu}), \quad n \rightarrow \infty.$$

Hence,

$$d_n = \frac{n^{\xi-1/2}}{5\pi} \int_0^1 \cos\left(\frac{2\pi n}{t}\right) \sqrt{t} dq(t) + O(n^{-\mu}), \quad n \rightarrow \infty. \quad (3.27)$$

The left-hand side of (3.27) does not depend on $\xi \in (1/2, 5/8)$. Therefore, writing (3.27) for some $\xi_1 > \xi$ from the same interval and then subtracting the previous equality, we obtain

$$n^{\xi_1-1/2} \left[1 - n^{\xi-\xi_1}\right] \int_0^1 \cos\left(\frac{2\pi n}{t}\right) \sqrt{t} dq(t) = O(n^{-\mu_1}), \quad n \rightarrow \infty$$

for some positive μ_1 . Hence,

$$\int_0^1 \cos\left(\frac{2\pi n}{t}\right) \sqrt{t} dq(t) = O(n^{1/2-\xi_1}), \quad n \rightarrow \infty$$

and (3.27) yields $d_n = O(n^{\xi-\xi_1}), \quad n \rightarrow \infty, \quad \xi < \xi_1$, completing the proof of Theorem 2. $\square$

Furthermore, the Salem-Zygmund theorem [18] shows that $d_n = o(1)$ implies that the Fourier-Stieltjes transform (1.1) $f(t) = o(1), |t| \rightarrow \infty$. Together with author's results in [16] it leads us to an immediate

Corollary 2. *Let $k \in \mathbb{N}$. Then the Fourier-Stieltjes transforms (1.1), (1.2) of the Minkowski question mark function and their consecutive derivatives $f^{(k)}(x), F^{(k)}(x)$ vanish at infinity.*

Finally, denoting by

$$d_{n,m} = \int_0^1 \cos(2\pi nt) dq^m(t), \quad m \in \mathbb{N} \quad (3.28)$$

the Fourier-Stieltjes coefficients of the power $q^m(t)$, we establish the following

Corollary 3. *We have $d_{n,m} = o(1), \quad n \rightarrow \infty$.*

Proof. Appealing to the principle of mathematical induction, we see that $d_{n,1} \equiv d_n = o(1), \quad n \rightarrow \infty$. Hence assuming that $d_{n,k} = o(1), \quad n \rightarrow \infty, \quad 1 \leq k \leq m$, we will prove that $d_{n,m+1} = o(1), \quad n \rightarrow \infty$. To do this, we recall (1.3) to have the equality

$$d_{n,m+1} = \sum_{k=1}^{m+1} (-1)^k \binom{m+1}{k} \int_0^1 \cos(2\pi nt) dq^k(1-t) = \sum_{k=1}^{m+1} (-1)^{k+1} \binom{m+1}{k} d_{n,k}. \quad (3.29)$$

Consequently, if m is odd, we find from the previous equality

$$d_{n,m+1} = \frac{1}{2} \sum_{k=1}^m (-1)^{k+1} \binom{m+1}{k} d_{n,k} = o(1), \quad n \rightarrow \infty.$$

When m is an even number, we write

$$2d_{n,m+2} - (m+2)d_{n,m+1} = \sum_{k=1}^m (-1)^{k+1} \binom{m+2}{k} d_{n,k} = o(1), \quad n \rightarrow \infty. \quad (3.30)$$

Meanwhile, since $m+3$ is odd

$$\sum_{k=1}^{m+2} (-1)^{k+1} \binom{m+3}{k} d_{n,k} = 0,$$

or

$$(m+3)d_{n,m+2} - (m+3)(m+2)d_{n,m+1} = \sum_{k=1}^m (-1)^{k+1} \binom{m+3}{k} d_{n,k} = o(1), \quad n \rightarrow \infty.$$

The latter equality means

$$d_{n,m+2} - (m+2)d_{n,m+1} = o(1), \quad n \rightarrow \infty.$$

Hence, with (3.30) it gives

$$(m+2)d_{n,m+1} = o(1), \quad n \rightarrow \infty$$

and therefore $d_{n,m+1} = o(1), \quad n \rightarrow \infty$.

□

Acknowledgments

The work was partially supported by CMUP (UID/MAT/00144/2013), which is funded by FCT (Portugal) with national (MEC) and European structural funds through the programs FEDER, under the partnership agreement PT2020.

REFERENCES

- [1] G. ALKAUSKAS, The Minkowski's $\varphi(x)$ function and Salem's problem, *C.R. Acad. Sci. Paris, Ser. I*, 350 (2012), 137-140.
- [2] A. DENJOY, Sur une fonction réelle de Minkowski, *J. Math. Pures Appl.* **17** (1938), 105–151 (in French).
- [3] O. S. IVAŠEV-MUSATOV, On Fourier - Stieltjes coefficients of singular functions, *Izv. Akad. Nauk SSSR, Ser. Mat.*, **20** (1956), 179– 196 (in Russian).
- [4] T. JORDAN, T. SAHLSTEN, Fourier transforms of Gibbs measures for the Gauss map, arXiv:1312.3619.
- [5] D. MENCHOFF, Sur l'unicité du développement trigonométrique, *Comptes Rendus*, **163** (1916), 433–436 (in French).
- [6] F. W. J. OLVER, D. W. LOZIER, R. F. BOISVERT, AND C. W. CLARK, EDITORS, *NIST Handbook of Mathematical Functions*, Cambridge University Press, New York (2010).
- [7] T. PERSSON, On a problem by R. Salem concerning Minkowski's question mark function, arXiv:1501.00876.
- [8] A.P. PRUDNIKOV, YU. A. BRYCHKOV AND O. I. MARICHEV, *Integrals and Series: Vol. 1: Elementary Functions*, Gordon and Breach, New York (1986); Vol. 2: *Special Functions*, Gordon and Breach, New York (1986).
- [9] R. SALEM, On some singular monotonic functions which are strictly increasing, *Trans. Amer. Math. Soc.*, **53** (3) (1943), 427-439.
- [10] R. SALEM, On monotonic functions whose spectrum is a Cantor set with constant ratio of dissection, *Proc Nat. Acad. Sc. USA*, **41** (1) (1955), 49–55.

- [11] R. SALEM, *Algebraic Numbers and Fourier Analysis*, Heath Math. Monographs, Boston (1963).
- [12] E.C. TITCHMARSH, *An Introduction to the Theory of Fourier Integrals*, Clarendon Press, Oxford (1937).
- [13] W. VAN ASSCHE, Minkowski's question mark function. In: Open Problems, Edited by C. Berg, *Integral Transf. Spec. Func.* **26** (2) (2015), 92- 95.
- [14] N. WIENER, A. WINTNER, Fourier-Stieltjes transforms and singular infinite convolutions, *Amer. J. Math.* **60** (3) (1938), 513–522.
- [15] S. YAKUBOVICH, Corrigendum to the Note "The Fourier-Stieltjes transform of the Minkowski's $\varphi(x)$ function and an affirmative solution to Salem's problem", [C.R. Acad. Sci. Paris, Ser. I, 349 (2011), 633-636], *C.R. Acad. Sci. Paris, Ser. I*, **350** (2012), 147.
- [16] S. YAKUBOVICH, On some Rajchman measures and equivalent Salem's problem, *Commun. Math. Anal.* , 14 (2013), N 1, 28-41.
- [17] S. YAKUBOVICH, The affirmative solution to Salem's problem revisited, arXiv:1501.00141.
- [18] A. ZYGMUND, *Trigonometric Series*, Vols. I, II. Cambridge University Press, Cambridge (1988).

SEMYON YAKUBOVICH, Department of Mathematics, Faculty of Sciences, University of Porto, Campo Alegre st., 687, 4169-007 Porto, Portugal. syakubov@fc.up.pt